



Paul J. Cagle, M.D.
ShoulderDoctorNYC.com

PECTORALIS MAJOR REPAIR POSTOPERATIVE PROTOCOL

0-2 Weeks

- Sling immobilization is used for 6 weeks - the sling is only removed for hygiene, showering and therapy
- Begin active motion of the elbow, wrist and hand
- Return visit to the office for a 2 week visit

2-6 Weeks

- Sling immobilization is used for 6 weeks - the sling is only removed for hygiene, showering and therapy
- Continue active range of motion of the elbow, wrist and hand
- Begin progressive shoulder passive and gentle active-assisted shoulder range of motion in all planes with no motion restrictions
- No shoulder strengthening exercises
- Return visit to the office for a 6 week visit

6-12 Weeks

- Discontinue sling immobilization
- Continue active range of motion of the elbow, wrist and hand
- Begin active shoulder range of motion in all planes with no motion restrictions - Progress to regain full range of motion
- No shoulder strengthening exercises.
- Return visit to the office for a 12 week visit

3-6 Months

- Continue active range of motion of elbow, wrist and hand
- Continue active shoulder range of motion in all planes with no motion restrictions and maintain full range of motion
- Begin Pectoralis, Deltoid, Rotator Cuff and Periscapular muscle strengthening exercises - Advance strengthening exercise as tolerated
- Return visit to the office for a 6 month visit

This protocol represents an overview of the standard postoperative course. Each patient's medical condition is unique, and thus, variations may be needed to tailor the rehabilitation protocol to optimize recovery.